





NARMS – EB 1999

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint


■ Slaughter n=1438 ■ Diagnostic n=85

NARMS – EB 1999


Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken


Ceftiofur




Ceftriaxone



Cephalothin



Chloramphenicol



⬇ Breakpoint





■ Slaughter n=1438

■ Diagnostic n=85

NARMS – EB 1999

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint


■ Slaughter n=1438 ■ Diagnostic n=85

*Breakpoint Not Established

NARMS – EB 1999

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint


■ Slaughter n=1438

■ Diagnostic n=85

NARMS – EB 1999

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint

⬇ Slaughter n=1438 ⬇ Diagnostic n=85